

Late Oligocene-Early Miocene transition: Paleoclimatic and paleoceanographic changes linked to a dynamic East Antarctic Ice Sheet, off the Wilkes Land margin.

Dimitris Evangelinos¹, Carlota Escutia¹, Ariadna Salabarnada¹, Adrián López-Quirós¹, Hans Nelson¹, Luis Valero², Johan Etourneau¹, Francisco J. Rodríguez-Tovar³, Peter Bijl⁴, Francisco J. Lobo¹ and Marga García¹

¹Instituto Andaluz de Ciencias de la Tierra, CSIC-Univ. de Granada, Av. de las Palmeras,4, 18100, Armilla, Spain. dimevangelinos@correo.ugr.es ;

²Laboratory of Paleomagnetism CCiTUB and CSIC. Institut de Ciències de la Terra Jaume Almera, Barcelona, Spain;

Universitat Autònoma de Barcelona, Facultat de Ciències, 08193 Bellaterra, Barcelona, Spain;

³Departamento de Estratigrafía y Paleontología, Facultad de Ciencias, Universidad de Granada, 18002 Granada, Spain.;

⁴Palaeoecology, Institute of Environmental Biology, Faculty of Science, Laboratory of Palaeobotany and Palynology, Utrecht University, Budapestlaan 4, 3584 CD Utrecht, The Netherlands.

Abstract

Low $\delta^{18}\text{O}$ values from deep-sea records during the Late Oligocene were attributed to deep-sea warming and decreased ice volume in Antarctica (e.g. Zachos et al., 2001). However, results from sedimentary records from the Antarctic continental margin indicate gradual Antarctic cooling during the Oligocene and large continental ice sheets on east Antarctica (e.g. Naish et al., 2001.) Resolving this controversy, Pekar et al. (2006) suggested expansion of warmer deep-waters from low latitudes, and reduction of deep-water production near the Antarctic continent. To provide new insights into the relationship between ice-sheet and oceanographic processes on a latitudinal transect across the Polar Front during the Oligocene-Miocene transition (OMT) we studied Late Oligocene- Miocene sediments from DSDP Leg 28 Site 269A (Hayes et al., 1975) and compare them with results from IODP Expedition 318 Site U1356 (Escutia et al., 2011; Salabarnada et al., in prep.) offshore the Wilkes Land margin. We use a combination of indicators in the sediment to help us reconstruct the paleoclimatic and paleoceanographic conditions during the late Oligocene to early Miocene. These include sedimentary facies and ichnological analyses, geochemical (XRF core-scanner) data, magnetic susceptibility, scanning electron microscope (SEM) images, and discrete mineralogical data (XRD). During the Late Oligocene to Early Miocene hemipelagic, downslope and along-slope processes influence the sedimentation in Site 269A. Reworking by bottom-currents is observed throughout the studied record showing changes in their strength and origin. During glacial times, glacial-marine deposits are winnowed by bottom currents, with preservation of laminations that contain coarse sand grains indicating increased current velocities. Turbidites and micro-slumps are scarce in this distal site protected from continental input by the Adélie Rift Block and could be linked to the extensive Mass Transport deposition during the Late Oligocene as recorded in U1356 (Escutia et al., 2011, Salabarnada et al., in prep.). Through the record, preservation of calcium carbonate is observed in cemented siltstone beds, and the high Ca-content is linked to the presence of calcareous microfossils (i.e. foraminifera and nannofossils). These cemented beds exhibit low MST values, high Si/Ti ratios associated with the presence of diatoms and radiolarians, and high Mn/Al ratios. All of these indicators point to interglacial periods with well-ventilated and productive waters. Towards the Oligocene-Miocene Boundary (OMB), Site 269A records an increase in Ca-rich intervals likely suggesting a prolonged expansion of warmer waters at this time. Our study on Site 269A and its comparison with Site U1356 suggests a very dynamic East Antarctic Ice sheet with several glacial/interglacial cycles and points to

various intrusion pulses of warmer bottom water from lower latitudes across the Late Oligocene to Early Miocene transition.

Keywords: Late Oligocene; East Antarctic Ice sheet; Oligocene-Miocene Boundary; Palaeoclimate

References

- Escutia, C., Brinkhuis, H., Klaus, A. and the Expedition 318 Scientists. 2011. Wilkes Land Glacial History: Cenozoic East Antarctic Ice Sheet evolution from Wilkes Land margin sediments. IODP Proceedings Volume 318 Integrated Ocean Drilling Program Management International.
- Hayes, D. E., Frakes, L. A., *et al.*, 1975. Initial Reports of the Deep Sea Drilling Project (28), U.S. Government Printing Office, Washington.
- Naish, T. R., Woolfe, K. J., Barrett, P. J., *et al.*, 2001. Orbitally induced oscillations in the East Antarctic ice sheet at the Oligocene/ Miocene boundary. *Nature*, (413), 719-723.
- Pekar, S. F., De Conto, R. M., Harwood, D. M., 2006. Resolving a late Oligocene conundrum: Deep-sea warming and Antarctic glaciation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, (231), 29-40.
- Salabarnada, A., Escutia, C., Nelson, H., Damuth, J., Brinkhuis, H., 2014. Varying depositional environments across the Oligocene-Miocene boundary and their relevance for East Antarctic ice sheet history: IODP Site U1356, Wilkes Land margin, in: *Geophysical Research Abstracts Vol. 16*, EGU2014-13705, 2014 EGU General Assembly 2014.
- Salabarnada, A., Escutia, C., Nelson, C.H., Roehl, U., Jimenez-Espejo, F., Evangelinos, D., Ikehara, M., McKay, R., Lopez, A., 2016. Obliquity paced contourite cyclicity in Antarctic sediments from the Wilkes Land (Site U1356) during Late Oligocene, in: *Geophysical Research Abstracts Vol. 18*, EGU2016-14854, 2016 EGU General Assembly 2016. p. 14854.
- Zachos, J., C., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms and aberrations in global climate 65 Ma to present. *Science*, (292), 686-693.