

## **Southern Ocean response to Antarctic glaciation in the mid Oligocene**

Claire E. Huck<sup>1</sup>, Myfanwy Wood<sup>1</sup>, Diederik Liebrand<sup>1</sup>, Ursula Röhl<sup>2</sup>, Paul Wilson<sup>1</sup>, Tina van de Flierdt<sup>3</sup> and Steven M. Bohaty<sup>1</sup>

<sup>1</sup>Ocean and Earth Science, National Oceanography Centre, University of Southampton, European Way, Southampton, SO14 3ZH UK, c.e.huck@soton.ac.uk;

<sup>2</sup>MARUM – Centre for Marine Environmental Sciences, University of Bremen, Leobener Strasse, D-3829, Germany;

<sup>3</sup>Department of Earth Science and Engineering, Imperial College London, Exhibition Road, London SW7 2AZ, UK.

### **Abstract**

The early stages of the modern 'Icehouse' climate state first developed in the Oligocene following rapid global cooling and the onset of Antarctic glaciation at the Eocene-Oligocene Transition (34 Ma). The size of the early Antarctic ice sheets, however, varied considerably, and a series of intense glacial-interglacial episodes occurred during the Mid Oligocene Glacial Interval (MOGI; ~28 to 26.3 Ma). In order to assess Southern Ocean paleoceanographic changes in response to this predominantly cool glacial phase, we investigated mid-Oligocene hemipelagic cores recovered at Ocean Drilling Program (ODP) Leg 113 Sites 689 and 690 on Maud Rise (eastern Weddell Sea). We generated new high-resolution x-ray fluorescence (XRF) scanning, benthic foraminiferal stable isotope, and fish-tooth neodymium (Nd) isotope records. The composited XRF records from these sites exhibit prominent orbital-scale cyclicity in iron/calcium (Fe/Ca) ratios (indicative of lithological changes clay vs. carbonate content), providing a record of carbonate dissolution intensity and/or changes in clay flux from the East Antarctic margin. Peak glacial conditions are inferred at ~27 Ma during long-term Eccentricity Cycle 67 based on the astronomically tuned benthic foraminiferal data from Site 689. During the peak glacial interval, a negative excursion in fish tooth Nd isotope record reveals a change in ocean circulation patterns, likely resulting from a shift in the relative contributions of deep-water sources to the Southern Ocean. This change in circulation is accompanied by a simultaneous increase in Fe/Ca ratios suggesting that enhanced weathering of nearby Antarctic bedrock by enhanced glacial activity may have also contributed to the short-term change in seawater chemistry during this time.

**Keywords:** Oligocene, Southern Ocean, Antarctica